arXiv:hep-ph/0203054v2 6 Apr 2002

# Effect of possible stronger neutrino interaction at $E_{\nu_\mu} \sim 10^{11}$ GeV upon the extraction of $\sin^2\theta_{\rm W}$ from the neutral-current cross section at NuTeV

Saul Barshay and Georg Kreyerhoff
III. Physikalisches Institut
RWTH Aachen
D-52056 Aachen
Germany

February 1, 2008

**Abstract**

The possibility exists that cosmic-ray neutrinos with energies of $\sim 10^{20}$ eV interact in the atmosphere with a cross section at the millibarn level, giving rise to some of the highest-energy air showers. In a specific dynamical model which can give rise to such a cross section, we show that there can be a small effect upon the extraction of the effective value of $\sin^2\theta_{\rm W}$ from $\nu_\mu$-hadron, neutral-current cross section data at NuTeV.

The possibility that neutrinos, with energies of the order of $10^{20}$ eV, have stronger interactions with hadrons than those which are given by electroweak theory, has been considered, specifically in connection with the puzzling observation of the highest energy cosmic-ray air shower events.[1, 2] Such energetic neutrinos can propagate from sources at cosmological distances (red shifts $\gtrsim 1$), undisturbed by the cosmic microwave background radiation, which photons largely prevent protons of $\gtrsim 0.6 \times 10^{20}$ eV from arriving here.[3, 4] Recently[5] we have explicitly calculated a $\sigma_{\nu_\mu}$ of $\sim 10^{-29}$ cm$^2$ at $E_{\nu_\mu} \sim 10^{20}$ eV, in a specific dynamical model which involves the hypothesis of a strong, parity-conserving interaction of $\nu_\mu$ with a hypothetical, point-like component of a $\pi^0$, leading to a very massive neutral lepton $L$, with $m_L \sim 2 \times 10^6$ GeV.$^{F1}$ The essential physical idea is that of neutrino structure at very small distances ($< 10^{-18}$ cm), due to hadron-like coupling to $L$. For this $m_L$, $E_{\nu_\mu}$ must reach $\sim 2 \times 10^{21}$ eV for production of real $L$. However, the amplitude for the diagram in Fig. (1), which involves a virtual $L$, results in a cross section above the electroweak $\sigma_{\nu_\mu} \sim 10^{-31}$ cm$^2$, already at $E_{\nu_\mu} \sim 10^{20}$ eV.[5] For a somewhat smaller $m_L \sim 0.3 \times 10^6$ GeV, the production of real $L$ is possible for $E_{\nu_\mu}$ above about $0.5 \times 10^{20}$ eV, which is close to where the GZK cut-off[3, 4] is located[1, 2] for protons coming from cosmological distances. In this case $\sigma_{\nu_\mu}$ can be of the order of millibarns at $E_{\nu_\mu} \sim 10^{20}$ eV. Then, a dip at the GZK cut-off, followed by a "bump-structure"[10] near to $10^{20}$ eV could occur in the air-shower

---

$^{F1}$ The possibility of an analogous hypothetical structure (with a charged pion) for the muon, can lead to a positive-definite, unusual contribution to $(g_\mu - 2)/2$ at the level of $10^{-9}$, for this value of $m_L$.[5, 6] However, in addition to the squared coupling, the magnitude is rather sensitive to possible damping at the structure vertex for virtual momentum near to $m_L$, where the integral gets its main contribution. At present, the significance of a possible experimental deviation[7] from expectation is reduced by the correction of a long-standing sign error in the calculation of a "standard" theoretical contribution.[8, 9] In this paper, we consider explicitly, only neutrino, effective neutral-current couplings.

energy spectrum, if some of the highest-energy events are induced by the primary neutrinos interacting in the atmosphere.[F2] Such a spectrum behavior is consistent with present data.[1, 2] This may only reflect insufficient exposure to obtain events at much higher energies. On-going and new cosmic-ray experiments will tell.[2]

The purpose of this paper is to show that the kind of hypothetical neutrino structure summarized above, can have an effect upon the extraction of $\sin^2\theta_{\mathrm{W}}$ from muon neutrino-hadron neutral-current cross section data at $E_\nu < 300$ GeV. The motivation for this is the recent NuTeV result[11] which indicates $\sin^2\theta_{\mathrm{W}} = 0.2276 \pm 0.0017$, about 2% higher than the value deduced mainly from the precision data at LEP, $\sin^2\theta_{\mathrm{W}} = 0.2226 \pm 0.0004$.[12] The NuTeV value of $\sin^2\theta_{\mathrm{W}}$ corresponds to a $W$ mass about 300 MeV below the direct measurements at LEP II and at the Tevatron.[12] It may well be that the NuTeV value of an effective $\sin^2\theta_{\mathrm{W}}$ relevant to $\nu_\mu$-hadron scattering is to be explained in terms of violation of isospin symmetry in the parton distributions of the nucleon, or in terms of an asymmetry in the strange quark sea.[13] It is common to speculate[15] about exchange of higher mass $Z'$ bosons (usually in the TeV range), whose amplitude interferes with that from $Z$ exchange. The new amplitude must be such as to give destructive interference. This reduces the neutral-current cross section.[F3] In a particular case[13], the $Z'$ coupling is required to conserve parity (i. e. no axial vector coupling). This is not the only way to obtain a parity-conserving, effective four-fermion interaction for $\nu_\mu$-hadron neutral-current scattering. A short-distance neutrino structure, of the kind discussed above, can also be characterized by such a parity-conserving, effective interaction, and can involve a very high particle mass. Our result below, suggests that for $m_L \sim 0.3 \times 10^6$ GeV an effective $\sin^2\theta_{\mathrm{W}}$ obtained from analysis of NuTeV data, might be up to $\sim 0.005$ larger than the expected electroweak value. Thus, a quantitative connection is made to the possibility of stronger neutrino interactions at the highest energies.

In Fig. (2), we illustrate schematically the reduction of the hypothetical primary dynamics at very high $\sqrt{s}$ approaching $m_L$, to an approximate, effective four-fermion interaction relevant for the NuTeV $\nu_\mu$-nucleon scattering data at $E_{\nu_\mu} <$ 300 GeV. The essential assumption is the following. The calculation of a $\nu_\mu$ cross section[5] for the hypothetical process in Fig. (1), involves using a typical hadronic total cross section to give an approximate strength for the sum of processes included in the square of the lower "vertex". This is a controlling factor in the size of the $\nu_\mu$ cross section. On the other hand, interference of the standard-model amplitude from Fig. (3), with an effective amplitude from Figs. (2), requires a degree of coherence for processes within the left-hand "black box" in Fig. (2a).[F4] We assume that this is achieved as illustrated in Fig. (2b). The axial-vector $A^0$ related to the $\pi^0$ in Fig. (1), propagates from the hypothetical (undamped) $\nu_\mu - L$ vertex to interaction with the target nucleon via a hard parton-parton collision; the result is a scattered nucleonic parton and an isoscalar, scalar "jet"[F5] (denoted by the exchanged dou-

---

[F2] There must be a sufficient flux of neutrinos near to $10^{20}$ eV, say $> 10^{-19}$ cm$^{-2}$ s$^{-1}$ sr$^{-1}$. A possible origin is from the two-body decay of a form of massive dark matter, with a lifetime ($\propto m_{\nu_\tau}^{-2}$)which is many orders of magnitude greater than the present age of the universe.[10] Neutrino-initiated air showers do not necessarily fall off strongly at larger zenith angles. (An indirect, dynamical mechanism which involves a sizable flux of very high-energy neutrinos as the origin of an air-shower "burst", postulates resonant interaction with CMB neutrinos (for non-zero mass), through a light $Z'$[13], as in ref. 14 for $Z$.)

[F3] This is the likely direction implied by the experimental effect in the measured ratio $(\sigma^{\mathrm{NC}}_{\nu_\mu}/\sigma^{\mathrm{CC}}_{\nu_\mu}) \sim 1/2 - \sin^2\theta_{\mathrm{W}} + (5/18)(\sin^2\theta_{\mathrm{W}})^2$, (given the $\sim 2$% precision in the measurements[16] of $\sigma^{\mathrm{CC}}_{\nu_\mu}$).

[F4] In Figs. (2,3), the box on the right represents hadronic fragmentation of partons, after the basic partonic collision process.

[F5] The jet is in effect, a neutral pair of pions, in the present context of hypothetical (point-like) $\pi^0$, $A^0$ interactions at the $\nu_\mu - L$ effective vertices.

ble line ($j^0$) in Figs. (2b,c)). In Fig. (2c), two vertices (with strength magnitude $|gg'| \sim g^2$) and the intermediate state on the neutrino line, characterized by the very high mass $m_L$, are brought together to an effective form $(g^2/m_L)(\overline{\nu}\gamma_\mu\gamma_5\nu)$. Similarly for the (coherent) intermediate states in the partonic "black box" on the quark line in Fig. (2b), with an assumed real part for the effective vertex. An effective Lagrangian can be approximately represented by the form

$$\mathcal{L}_{\rm eff} = -\left(\frac{\sqrt{2}g^2}{m_L m}\right)\left(\overline{\nu_\mu}\gamma_\mu\gamma_5\nu_\mu\right)\left(\overline{q}\gamma^\mu\gamma_5\tau_3 q\right) \tag{1}$$

where $q$ represents a $u$ or $d$ quark, and $m$ is an effective mass given by the product of significant numerical factors of $\pi$ (see Eq. (4) below) and a dynamical mass $\sim 3$ GeV which characterizes the "black box". The negative sign before $\mathcal{L}_{\rm eff}$ ( for $q = u$) corresponds to destructive interference of the amplitude with the standard-model amplitude from Fig. (3). The negative sign can arise from the structure of the intermediate states in Fig. (2) and the overall sign of the product of effective vertices. This sign appears to arise naturally when the hadronic black box in Fig. (2b) involves a hard parton scattering (gluon exchange). The order of magnitude of the ratio of the amplitude from Fig. (2c) to that from Fig. (3) is

$$|R| \sim \left(\frac{g^2}{e^2}\right)\left(\frac{m_Z}{m_L}\right)\left(\frac{m_Z}{m}\right) \sim 0.002 \tag{2}$$

for $g^2/(4\pi) \sim 1$, $m_L \sim 0.3 \times 10^6$ GeV, $e^2/(4\pi) \cong \frac{1}{137}$, $m_Z \cong 90$ GeV and with $m \sim (20\pi^3)(3 \text{ GeV}) \sim 1.8 \times 10^3$ GeV (calculated in Eq. (4) below).

In Eq. (1), the effective strength is

$$\left(\frac{\sqrt{2}g^2}{m_L m}\right) \sim 0.33 \times 10^{-7} \approx \left(\frac{G_F}{\sqrt{2}}\right)\frac{(0.01)}{2} \tag{3}$$

The value $\sim (G_F/\sqrt{2})(0.01)$ has been deduced [13] phenomenologically as the approximate necessary strength of an effective four-fermion interaction which is, a priori, required to be $SU(2)_L$ invariant (we use only a corresponding parity-conserving, axial-axial term involving $\overline{\nu}\nu$ and $\overline{u}u(\text{-}\overline{d}d)$). In the present model, the effective strength in Eq. (3) is directly fixed by the hypothesis of relatively strong, parity-conserving neutrino interactions (thus $g^2/4\pi \sim 1$ compared to $e^2/4\pi$), at $E_\nu \sim 10^{20}$ eV (corresponding to $m_L \lesssim 10^6$ GeV). The effective mass $m$ is determined by the dynamical structure of the model, as we now estimate. The neutrino cross section (here assumed to be relevant for p or n in an isoscalar target), calculated approximately from the amplitude in Fig. (2b), is

$$\begin{aligned}
\Delta\sigma_{\nu_\mu} \quad &\sim \quad (g^2)^2\left\{\left(\frac{1.22}{(4\pi)^2}\right)^2\left(\frac{\langle n_j\rangle\sigma_j}{2\pi^2}\right)\right\}\left\{\left(\frac{4}{3\pi}\right)\left(\frac{s}{4m_L^2}\right)\right\} \\
&= \quad (g^2)^2\left(\frac{1}{m^2}\right)\left\{\left(\frac{4}{3\pi}\right)\left(\frac{s}{4m_L^2}\right)\right\} \cong (8\pi)^2\left\{\frac{1}{(1.8\text{ TeV})^2}\right\}(4.7\times 10^{-10}) \\
&\cong \quad (0.78\times 10^{-31}\text{ cm}^2)(4.7\times 10^{-10}) \cong 0.37\times 10^{-40}\text{ cm}^2
\end{aligned} \tag{4}$$

In order to make clear the origin of the magnitude in this dynamical model, three separate factors are enumerated in detail in Eq. (4). There is the overall strong strength from the two interactions on the neutrino line, represented by the magnitude $|gg'| \sim g^2$ ($g^2/4\pi \sim 2$, is used, as in ref. 5). The second factor in curly brackets ($\{\ldots\}$) is kinematical, essentially the phase space. At $\sqrt{s} \sim 20$ GeV, used in Eq. (4) for illustration, this is a small number of the order of $10^{-9}$. This number is controlled by the large mass $m_L \sim 0.3 \times 10^6$ GeV, whose size is approximately fixed by the possibility of strong neutrino interactions at extremely high energy, $\sim 10^{11}$ GeV. The first factor in curly brackets represents the (inverse) square of an effective mass $m \sim (20\pi^3)(3 \text{ GeV})$; this controls the effective strength of the hadronic "vertex". The mass has a large kinematic factor ($\sim 20\pi^3$) which is related to the specific dynamics, i. e. to the integration over virtual momenta, and to the approximate parameterization of the "black box" in Fig. (2b), in terms of a measured cross section for a hard scattering process. For the latter we use a typical jet cross section at relatively[17] low $\sqrt{s}$, $\sigma_j \sim 2 \times 10^{-29}$ cm$^2$ in a central rapidity interval $|\Delta y| \sim (1 - (-1)) = 2$. We use this as a (fixed) parameter only in a limited range of the intergration momentum[F6], $\sim 2$ GeV to $\sim 3.7$ GeV; $\sigma_j$ falls away at high loop momentum.[F7] With the jet multiplicity $\langle n_j \rangle \cong 2$, the quantity $(\langle n_j \rangle \sigma_j)^{-1/2}$ translates into a mass of $\sim 3$ GeV, which can be considered as the intrinsic, mass parameter characteristic of the "black box" in Fig. (2b).

Using the estimate for $\Delta\sigma_{\nu_\mu}$ in Eq. (4), and the empirical $\nu_\mu$ charged-current cross section[16], $\sigma_{\nu_\mu}^{\text{CC}}(E_{\nu_\mu} \sim 200 \text{ GeV}) \cong 1.36 \times 10^{-36}$ cm$^2$, we calculate the magnitude of the interference term between the amplitudes from Figs. (2,3), in ratio to $\sigma_{\nu_\mu}^{\text{CC}}$,

$$|I| \cong 2\sqrt{\frac{\Delta\sigma_{\nu_\mu}}{\sigma_{\nu_\mu}^{\text{CC}}}}\sqrt{\frac{\sigma_{\nu_\mu}^{\text{NC}}}{\sigma_{\nu_\mu}^{\text{CC}}}} \cong 2(0.0052)(0.55) = 0.0057 \tag{5}$$

This is approximately independent of the specific $E_{\nu_\mu}$ used above for illustrating the numbers. In Eq. (5), we use $\sim 0.29$ for the ratio of neutral to charged-current cross sections (i. e. as calculated[F3] for $(\sin^2\theta_{\text{W}})_{\text{LEP}} = 0.223$). A negative interference term of this magnitude corresponds to an effective value of $\sin^2\theta_{\text{W}}$ in the NuTeV experiment which is about 0.005 larger than the (electroweak) value determined from the LEP measurements.

From $\overline{p}p \to l^+l^- + X$ interactions, CDF has set limits[19] on quantities analogous to $\epsilon \sim 0.01$ in Eq. (3), but referring to effective four-fermion interactions involving quarks and a $\mu^-\mu^+$ (or $e^-e^+$) pair, (approximately, the $|\epsilon|$ are[13]$\lesssim 0.04$). It is noteworthy that the hypothetical, specific $\nu_\mu$ structure in Fig. (2) does not have a counterpart for $\mu^-$. This is because the scalar "jet" is neutral[F5]. Thus, the experimental limits[19] from $q\overline{q} \to \mu^-\mu^+$ are not directly relevant here (as they are for phenomenological interaction forms utilized in ref. 13).

There is a potentially interesting "charge" radius effect for $\nu_\mu$, implied by the hypothetical structure. If $\rho^0$-exchange can occur in place of the $A^0$ in Fig. (2c), a charge radius is generated by replacing the lower vertex by that for $\gamma \to \rho^0\pi^+\pi^-$, with a magnitude estimated as

$$|\langle r^2 \rangle_{\nu_\mu}| \sim \left(\frac{g^2}{64\pi^2}\right)\left(\frac{1}{\mu m_L}\right) \sim 1.6 \times 10^{-35} \text{ cm}^2 \tag{6}$$

The mass parameter $\mu \sim 264$ MeV, is estimated from the measured partial width[16] $\Gamma(\rho^0 \to \pi^+\pi^-\gamma) \sim 1.5$ MeV. Empirically[16], $|\langle r^2 \rangle_{\nu_\mu}| \lesssim 10^{-32}$ cm$^2$. More generally, this represents a limit on certain non-standard contributions to $\nu_\mu$ scattering[16]; $\Delta\sigma_{\nu_\mu}$ in Eq. (4) is much less. At the Z-pole, a similar "charge" radius can affect, in principle, the magnitude of the axial-vector (and possibly, vector) effective coupling

---

[F6] Note that a corresponding typical cross section[18] for $\pi^0$, at low $\sqrt{s}$ in this limited range for transverse momenta, is comparable, about $0.4 \times 10^{-30}$ cm$^2$. It falls away more rapidly than does $\sigma_j$, at higher transverse momenta.

[F7] The approximate integration over virtual momenta $Q$, with $\sigma_j$ fixed over a limited range, involves a factor $\ln(Q^2_{\text{max}}/Q^2_{\text{min}})$. This is of order unity; we use $2\ln(3.7/2) \sim 1.22$. If taken as unity, the interference term in Eq. (5) has magnitude $\sim 0.0047$.

to $\nu_\mu \overline{\nu_\mu}$. We have, $m_Z^2 |\langle r^2 \rangle_{\nu_\mu}| \sim 0.03\%$, which corresponds to a change in this partial width of about ±0.1 MeV. This means that the effective number of light neutrinos can be less than three[15]. There is $\sim 3\%$ accuracy in the measurement[20] of the $\nu_\mu$-coupling to $Z$ from $\nu_\mu$-electron scattering. The direct measurement[21] of the invisible decay width of the $Z$, is also with $\sim 3\%$ accuracy. There is a fit value[16] with $\sim 0.3\%$ uncertainty.

We summarize the model, the essential hypothesis for which involves the neutrino production by effectively strong interactions, of a very massive neutral lepton at the highest energies. The dynamics involves hypothetical, strong interactions of point-like components of exchanged hadronic entities which couple a neutrino to the heavy lepton $L$; thus, a corresponding effective neutrino substructure at very short distances. The resulting calculated cross section[5] can reach above the millibarn level at $E_\nu \sim 10^{20}$ eV. The enhanced cross section occurs already below threshold[5], and increases through the threshold for production of real $L$. Clearly, the exchange mechanism in Fig. 1 involves all angular momenta (this is not a unitarity-bounded, S-wave effect, as illustrated in [23]). In accord with estimates in [23], the scale (and dimension) of the new cross section is set by a hadronic total cross section (from the square of the lower "vertex" in Fig. 1; as given explicitly in Eq. (5) in [5]). In this paper, we have "extrapolated" from the hypothetical high-energy dynamics, to a small effect at NuTeV energies. This arises from dynamical and kinematical factors which are estimated numerically, and which are explicitly enumerated and explained in the context of the high-energy model, following Eq. (4) above. At NuTeV $\sqrt{s}$, the relevant effective mass parameter is calculated to be about 4.7 TeV.

In conclusion, we note that while the present NuTeV effect might eventually get an explanation in terms of asymmetries in parton distributions[13], ideas concerning hypothetical new particles[15, 13] are not necessarily restricted to masses below a few TeV. In the effective four-fermion interaction generated by structure, in Eq. (1), while the overall effective mass parameter is estimated as $\sim \sqrt{m m_L / g^2} \sim 4.7$ TeV, the particle mass $m_L \sim 300$ TeV, is much higher. This is because the hypothesis is made of a dynamical connection between the possibility of stronger neutrino interaction in the atmosphere, reaching of the order of millibarn cross sections at the highest energies $\sim 10^{11}$ GeV, and a small effect upon $\nu_\mu$-hadron scattering below a few hundred GeV. This relates two presently puzzling phenomena: the cosmic-ray air showers at $10^{20}$ eV and the possible anomalous deviation in the $\sin^2\theta_{\mathrm{W}}$ extracted from NuTeV data.[F8]

## Added note

After completion of this work, authors have called our attention to recent work on large cross sections from brane theory; we include some references[24, 25, 26]. See also [27] and [28], and the earliest related work [29] which we are aware of.

## References

[1] M. Takeda et al., Phys. Rev. Lett. **81**, (1998) 1163

[2] M. Nagano and A. A. Watson, Rev. Mod. Phys. **72** (2000) 689

---

[F8] A limit on the possibility of a small amplitude arising from a hypothetical, point-like component of the pion coupling, to vector bosons[22], is discussed in the conclusion of[5]. Possible observational consequences are: (1) a partial width for $Z^0 \to \pi^+\pi^-$ of the order of an MeV, and (2) a quite small, but negative value, for the $S$ parameter[15], corresponding to a downward shift in the $Z$ mass of the order of MeV.

[3] K. Greisen, Phys. Rev. Lett. **16** (1966) 748

[4] G. T. Zatsepin nad V. Kuzmin, JETP Lett. **4** (1966) 78

[5] S. Barshay and G. Kreyerhoff, Eur. Phys. J. **C23** (2002) 191

[6] B. E. Lautrup, A. Peterman and E. de Rafael, Phys. Rep. **3C** (1972) 193

[7] Muon $(g_\mu - 2)$ Collab., H. M. Brown et al., Phys. Rev. Lett. **86** (2001) 2227

[8] M. Hayakawa and T. Kinoshita, hep-ph/0112102

[9] M. Knecht and A. Nyffeler, hep-ph/0111058

[10] S. Barshay and G. Kreyerhoff, Nuovo Cimento **112A** (1999) 1463; **112A** (1999) 1469

[11] G. P. Zeller et al., hep-ex/0110059v2

[12] J. Drees, hep-ex/0110077v1

[13] S. Davidson et al., hep-ph/0112302v2

[14] T. Weiler, Phys. Rev. Lett. **49** (1982) 234 and erratum ibidem **12** (2000) 379

[15] J. Erler and P. Langacker, Phys. Lett. **B456** (1999) 68
J. Erler and P. Langacker, Phys. Rev. Lett. **84** (2000) 212
P. Langacker, hep-ph/0102085

[16] Particle Data Group, Rev. Part. Phys., Eur. Phys. J. **C15** (2000) 379

[17] AFS Collab., T. Akesson et al., Phys. Lett. **123B** (1983) 133

[18] C. Kourkoumelis et al., Z. Phys. **C5** (1980) 95

[19] CDF Collab., F. Abe et al., Phys. Rev. Lett. **79** (1997) 2198

[20] CHARM II Collab., P. Vilain et al., Phys. Lett. **B320** (1994) 203

[21] L3 Collab., M. Acciarri et al., Phys. Lett. **431** (1998) 199

[22] S. Barshay, Phys. Lett. **B286** (1992) 375

[23] G. Burdman, F. Halzen, and R. Gandhi, Phys. Lett. **B417** (1997) 107

[24] L. A. Anchordoqui, J. L. Feng, and H. Goldberg, hep-ph/0202124

[25] E-J. Ahn, M. Cavaglià, and A. V. Olinto, hep-th/0201042

[26] P. Jain, S. Kar, S. Panda, and J. P. Ralston, hep-ph/0201232

[27] G. Domokos and S. Kovesi-Domokos, Phys. Rev. Lett. **82** (1999) 1366

[28] J. Bordes et al., Astropart. Phys. **8** (1998) 135

[29] V. Berezinsky and G. Zatsepin, Phys. Lett. **B28**, (1969) 423

## Figures



Figure 1: Feynman diagram for a contribution to the interaction cross section of an extremely high energy, cosmic-ray neutrino with an atmospheric nucleon, mediated by a virtual $L^0$.



Figure 2: For $\nu_\mu$-hadron scattering at $\sqrt{s} \sim 20$ GeV, a schematic sequence leading to the approximate effective four-fermion interaction in Eq. (1), as described in the text and footnotes F4-7. The effective vertices on the neutrino line are assumed to be point-like, up to momenta near to the scale of $m_L$. The loop integration momentum is taken as relevant for jet production, but only a limited range is used in estimating the amplitude for Eq. (4), 2 GeV $\lesssim$ |Q| $\lesssim$ 3.7 GeV, because $\sigma_j$ falls away at high $|Q|$.



Figure 3: The standard model interaction.